CMK2012-016-016

CMK2012-016-016

CMK2005-210-007 x LADYANNA

MEIJER POTATO
Building on potential

Purpose: French fries

Characteristics:

Maturity	5,9	
Colour of skin	Y	
Colour of flesh	6,3	
Dry matter percentage	22,4	
Suitability for cooking	7,9 B	A B C D
Shape of tuber	LO	
Tuber length (50-60 mm)	8,6 cm	
Average size (mm)	57	
Dormancy	7,7	

Added Value:

Uniform tuber size in high marketable yields.

Robust plant, tolerant to climatic stress.

Excellent frying colour throughout the storage period.

Gives high yields from all soil types.

Long dormancy suitable for long storage.

Resistances :

Common scab	7
Bruise index (0-50)	11
Secondary growth	7
Yntn-virus	10
Foliage blight	4
Fusarium	7

Ro1	9		
Ro2/3	8	wz 1 (D1)	5
Ro5		wz 2 (G1)	
Pa2	7	wz 6 (O1)	3
Pa3	1	wz 18 (T1)	5

Planting distance:

Size	Rowspace (cm between plants) 75 cm	90 cm	Plants/hectare
28/35	33	28	40404
35/50	41	34	32520
50/55	44	37	30303

Meijer Potato will not be responsible for possible negative effects originating from this advice.

CMK2012-016-016

MEIJER POTATO
Building on potential

Purpose: French fries



Planting

Wait for eyes waking up before planting.
Do not plant cold seed in cold soil.
Don't de-sprout the seeds before planting.
Plant at a normal depth.
Seems to be have a susceptibility to powdery scab; choose the right soils.

Fertiliser input (kg/ha):

Depending on the soil supply and applicable national legislation; total amount required by the variety.



			Total
Nitrogen:	(N)	250	**250**
Phosphate:	(P_2O_5)	110	**110**
Potassium:	(K_2O)	350	**350**

To prevent mineral deficiencies, apply these elements in time (B, Mg, Mn, S)

Variety with low needs of nitrogen.
Too much nitrogen affect the yield in a negativ way.

Crop protection

Susceptible to the use of 150 g/ha of Metribuzin at post-emergence.
Variety susceptible to foliar late blight, do not neglect blight control.
Start late blight control in time.

Harvest and storage

Wait for complete seed disintegration before harvesting.
After harvest, start immediately with drying the potatoes.

Storage:
7 °C
8 months

Key advice to succeed this variety:

Wait for eyes waking up before planting.
Seems to be have a susceptibility to powdery scab; choose the right soils.
Susceptible to the use of 150 g/ha of Metribuzin at post-emergence.
Variety susceptible to foliar late blight, do not neglect blight control.
After harvest, start immediately with drying the potatoes.

Meijer Potato will not be responsible for possible negative effects originating from this advice.

21-12-2020